

**Clinical Management Plan for Radiotherapy related Toxicity: For Pelvic/Abdominal
Radiotherapy for Gynaecological Cancers**

Name of Patient/ID: <i>Attach patient sticker if available</i>		Sensitivities/allergies:	
Independent Prescriber(s): Contact details:		Supplementary Prescriber(s) Alison Snook, Sarah Griffiths, Pauline Humphrey, Jancis Kinsman Hazel Boyce Helen Cullimore Contact details:	
Condition(s) to be treated Diagnosis: Radiotherapy course/schedule: Radical/Adjuvant/Palliative: Concurrent Chemo:		Aim of treatment: Control of radiotherapy related toxicity	
Medicines that may be prescribed by SP:			
Preparation	Indication	Dose schedule	Specific indications for change in management/ referral back to the IP
Loperamide Hydrochloride	Onset of radiation induced diarrhoea RTOG grade 1 and 2	As detailed in Section 1.4.2 BNF	Diarrhoea not controlled with 16mg Loperamide daily. Evidence of onset of dehydration. Development of RTOG grade 3 toxicity.
Scheriproct Ointment	Anal/perianal skin reaction or soreness due to diarrhoea	As detailed in Section 1.7.2 BNF	Significant rectal bleeding
Fluconazole	Vaginal Candidiasis	A single dose of 150mg, as detailed in Section 5.2 BNF	If no benefit or symptomatically worse
Trimethoprim	Uncomplicated lower urinary tract infection – symptomatic and positive dipstick test or positive microscopy	3 day course, as detailed in Section 5.1 BNF	Microscopy sensitivity requires another antibiotic. Symptoms not resolving after course completed.

BRISTOL HEAMATOLOGY AND ONCOLOGY CENTRE

Metoclopramide	Radiation induced nausea	As detailed in Section 4.6 BNF	If no benefit or symptomatically worse	
Domperidone	Radiation induced nausea	As detailed in Section 4.6 BNF	If no benefit or symptomatically worse	
Ondansetron	Radiation induced nausea	As detailed in Section 4.6 BNF	If no benefit or symptomatically worse	
Paracetamol	Pain as per WHO Pain Ladder Step 1a	As detailed in Section 4.7.1 BNF	Pain not controlled	
Ibuprofen	Pain as per WHO Pain Ladder Step 1b	As detailed in Section 10.1.1 BNF	Pain not controlled	
Co-codamol 30/500	Pain as per WHO Pain Ladder Step 2	As detailed in Section 4.7.1 BNF	Pain not controlled	
<p>Guidelines or protocols supporting Clinical Management Plan: ASCO guidelines: Recommended Guidelines for the Treatment of Cancer Treatment-Induced Diarrhoea. Journal of Clinical Oncology; 22 (14): 2918-2026 WHO Pain Ladder ASWCS Network Anti-emetic Policy (Version 2006.1) BHOC Radiotherapy Protocol for Gynaecological Cancers UH Bristol Foundation Trust Medicines Code: Non-medical prescribing policy UH Bristol Foundation Trust Medical division Empirical Antibiotic Guidelines</p>				
Frequency of review and monitoring by:				
Supplementary Prescriber Weekly or sooner if patient reports problems		Supplementary Prescriber and Independent Prescriber At request of Supplementary Prescriber or Patient's request		
<p>Process for reporting ADRs: Yellow Card system in BNF UH Bristol Foundation Trust Clinical Incident Form as per Medicine Management Policy Documentation in medical notes and radiotherapy treatment sheet Inform IP</p>				
<p>Shared record to be used by IP and SP: Radiotherapy Treatment Sheet and Oncology notes Patient Held Records (when available)</p>				
Agreed by Independent Prescriber(s)	Date	Agreed by Supplementary Prescriber(s)	Date	Date agreed with Patient/carer